arXiv:hep-lat/0110226v1 31 Oct 2001

Vector condensation in QCD

K. Splittorff[a*]

[a]The Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

The properties of the QCD vacuum at quark chemical potentials on the order of the pion mass and at zero temperature are by now quite well understood. The insight into this non-perturbative regime of QCD is obtained using chiral perturbation theory at non-zero quark chemical potentials [1–8]. The results apply not only to ordinary QCD but also to two color QCD and to QCD with quarks in the adjoint representation of the color group. Some of these realizations of QCD does not have a sign problem at finite chemical potential and hence can be studied by standard lattice methods as well. This has lead to a fruitful interplay between lattice calculations and chiral perturbation theory (see [6] for an extensive list of references). The central observation of all these studies is the condensation of specific goldstone channels. The critical chemical potentials at which the condensation occur can be understood in terms of the Bolzman weight in the partition function since the effective theory at lowest order describes a non-interacting Bose gas. For example, non-zero densities in QCD with $N_c = N_f = 2$ set in for isospin chemical potential $\mu_I > m_\pi$ or baryon chemical potential $\mu_B > m_\pi$ [3]. This is so since in this theory there is no goldstone mode with non-zero third component of isospin and non-zero baryon charge.

The effects at next to leading order in the chiral expansion have been studied in two colour QCD at finite baryon chemical potential [7]. The renormalized theory at next to leading order confirms that the critical baryon chemical potential is $\mu_B = m_\pi$. In addition the critical exponents at the phase transition was calculated. Surprisingly, they remain at the mean field values.

The inclusion of a set of massive spin-one vectors, A_ν, in lowest order chiral perturbation theory for two-color QCD was studied in [9]. In this case the effective Lagrangian must respect global $SU(2N_f)$ transformations and is given by

$$\begin{aligned}\mathcal{L}_{eff} &= f^2 \,\mathrm{Tr}\left[D_\nu \Sigma D^\nu \Sigma^\dagger\right] + m_V^2 \,\mathrm{Tr}\left[A_\nu A^\nu\right] + h^2 \,\mathrm{Tr}\left[A_\nu \Sigma A^{T\nu} \Sigma^\dagger\right] \\ &+ i\,s f^2 \,\mathrm{Tr}\left[A_\nu \Sigma D^\nu \Sigma^\dagger\right] + f^2 m_\pi^2 \mathrm{Tr}[\mathcal{M}\Sigma + \mathcal{M}^\dagger \Sigma^\dagger] - \frac{1}{2g^2}\mathrm{Tr}\left[F_{\rho\nu} F^{\rho\nu}\right] \ , \end{aligned} \tag{1}$$

where $\Sigma \in SU(2N_f)/Sp(2N_f)$, $A_\nu \in SU(2N_f)$, and $F^{\rho\nu} = \partial^\rho A^\nu - \partial^\nu A^\rho - i\left[A^\rho, A^\nu\right]$. The effect of a non-zero baryon chemical potential, μ_B, in this theory was examined in [6]. The chemical potential enters through (B is the baryon charge matrix)

$$\partial_0 \Sigma \to \partial_0 \Sigma - i\mu_B \left[B, \Sigma\right] \quad \text{and} \quad \partial_0 A_\rho \to \partial_0 A_\rho - i\mu_B \left[B, A_\rho\right] \ . \tag{2}$$

*In collaboration with J.T. Lenaghan, F Sannino, J.J.M. Verbaarschot, D. Toublan, D.T. Son, and M.A. Stephanov



Figure 1. The masses of the vector states in $N_c = N_f = 2$ QCD. The masses on the left are triply degenerate. At $\mu_B = 0$ the vector mass splitting between vector channels associated with the broken, X, and unbroken, S, generators is $M_X^2 - M_S^2 = 2g^2 f(h + 2 - s)$. The parameters in the effective Lagrangian are taken as $(h, s, m_V, f, g) = (0, 0, 4m_\pi, m_\pi, 1)$.

From the quadratic terms in A_ν of $\mathcal{L}_{eff}$ one can calculate the masses of the $(2N_f)^2 - 1$ vectors as a function of μ_B. An explicit calculation for $N_f = 2$ shows that [6] three of the 15 vector masses are degenerate and independent of μ_B. The latter 12 vector states mix as a function of μ_B. The mixing is not complete, since it splits into 4 sections each containing 3 vector states. Three of these sections have the same mass spectrum. Moreover all 4 sectors have a massless mode at $\mu_B = M_S$, where M_S is the mass of the lightest vector at $\mu_B = 0$. The fact that the vector mass vanishes suggests that these 4 modes condense at this and higher values of the chemical potential.

The method outlined here and given in detail in [6] directly applies to three color QCD as well as to QCD with quarks in the adjoint representation. For instance, one can calculate the vector masses in three colour QCD at finite isospin chemical potential, μ_I, where condensation in the ρ-channel is expected to occur at $\mu_I = m_\rho$. Note that this prediction can be examined by means of lattice simulations.

REFERENCES

1. R. Rapp, T. Schaefer, E.V. Shuryak, M. Velkovsky, Phys.Rev.Lett. **81** (1998) 53.
2. J.B. Kogut, M.A. Stephanov, D. Toublan, Phys.Lett. **B464** (1999)183; J.B. Kogut, M.A. Stephanov, D. Toublan, J.J.M. Verbaarschot, A. Zhitnitsky, Nucl.Phys. **B582** (2000) 477.
3. D.T. Son, M.A. Stephanov, Phys.Rev.Lett. **86** (2001) 592.
4. K. Splittorff, D.T. Son, M.A. Stephanov, Phys.Rev.**D64** (2001) 016003.
5. J.B. Kogut, D. Toublan, Phys.Rev. **D64** (2001) 034007.
6. J. T. Lenaghan, F. Sannino, K. Splittorff, `hep-ph/0107099`.
7. K. Splittorff, D. Toublan, J.J.M. Verbaarschot, `hep-ph/0108040`.
8. J.B. Kogut, D. Toublan, D.K. Sinclair, Phys.Lett. **B514** (2001) 77.
9. Z. Duan, P.S. Rodrigues da Silva, F. Sannino, Nucl.Phys. **B592** (2001) 371.